Ouachita Baptist University

Scholarly Commons @ Ouachita

Honors Theses

Carl Goodson Honors Program

5-1974

Response of Insects to U.V. Light and Varying Intensities of White Light

Tim Matthews Ouachita Baptist University

Follow this and additional works at: https://scholarlycommons.obu.edu/honors_theses



Part of the Entomology Commons

Recommended Citation

Matthews, Tim, "Response of Insects to U.V. Light and Varying Intensities of White Light" (1974). Honors Theses. 549.

https://scholarlycommons.obu.edu/honors_theses/549

This Thesis is brought to you for free and open access by the Carl Goodson Honors Program at Scholarly Commons @ Ouachita. It has been accepted for inclusion in Honors Theses by an authorized administrator of Scholarly Commons @ Ouachita. For more information, please contact mortensona@obu.edu.

V. ANALYSIS OF RESULTS

Although theoretically insects react more strongly to ultraviolet light than to ordinary white light, such was not the case in this experiment. There are several reasons for this deviation from the expected. One is the fact that there were other lights (within the range of the insects' vision) of greater intensity than those lights used in the experiment. This would cause the insects to react towards the more intense light source instead of the collector light. Although this reinforces the previously stated theory of "the greater the intensity, the greater the response", it voids the results concerning the insects reaction to ultraviolet light.

Ultraviolet light will draw more response than a white light of corresponding intensity, but there is a point at which the intensity will over-rule the response to the ultraviolet light.

Another factor that may have led to incorrect results is that of the intensity of the moon-light. On cloudy nights the reponse was greater than the response illicited on clear nights.

The temperature also had a very definite effect on the results. The lower the temperature the less the response. This relationship between temperature and response is an indirect one rather than a direct one. It is most likely that the temperature either killed or slowed down the insects, and did not actually effect the electro-chemical response.

In order to obtain valid results one would need to maintain strict controls on the temperature, wind currents, surroundings and humidity. The method of using the numbers of insects collected as the basis for experimental conclusions about the response of insects to variations of intensity and wave-length is not actually a valid method. A more precise method would be to record (if possible) the electrical responses of a single visual cell of the compound eye.


RESPONSE OF INSECTS TO U.V. LIGHT AND VARYING INTENSITIES OF WHITE LIGHT

- I. OBJECT: To determine the sensitivity of insects to ultraviolet light and also to verify their response to varying intensities of white light.
- II. THEORY: Insects are generally sensitive to light of any wave-length. Some insects' reactions are due to the response of specialized epidermal cells whereas the center of most response is in the compound eye. The reaction is generally thought to be due to an electro-chemical reaction within the light sensitive cells of the compound eye, i.e. mainly the cells of the crystalline cone and the rhabdomere. Insects tend to react more strongly to light in the ultraviolet region although the mechanism(s) which cause this reaction have not been established.


It has been observed experimentally that the magnitude of the insects' response to white light is directly proportional to the intensity of the light. The actual cause of this response has also not been determined.

III. APPARATUS: Diagrams of the apparatus used are on the following page.

1. Box collector - for white light



2. Funnel collector



IV. DATA
FALL 1973

A. Box 250 Watt - white light

DATE	CATCH	TIME	REMARKS
10/9/73	5	2"	Weather- clear, cool
#1	4	5 "	
11	5	15"	
ti	4	30 "	•
10/11/73	7	2"	Weather- clear, cool, humid
11	4	5 "	
Ħ	5	15"	
, , , , , , , , , , , , , , , , , , , 	6	30 "	
10/15/73	4	5"	Weather- cool, humid
ti	3	5"	Began using only 5" exposure
11	4	5"	time.
10/19/73	3	5"	Damp. cooler (below 60°F)
11	2	5"	
tf	4	5"	
10/25/73	2	5"	Too cool for any reasonable re-
n	3	5 "	sponse.
11	2	5 "	
11/3/73	1	5"	Weather cool
11	_	5 "	
91	- :.	5"	
11/20/73	-	5"	Weather cool. Cold spell last
Ħ	-	5 "	week.
11		5"	

B. Box 100 Watt - white light

DATE	CATCH	TIME	REMARKS
10/9/73	4	2"	Weather clear, cool.
91	4	5."	•
tf	3	15"	
11	5	30"	
10/11/73	3	2"	Weather - clear, cool, humid
t1	2	5"	
11	5	15"	
15	3	30"	
10/15/73	3	5"	Almost all of the catch was
f †	2	5 "	mosquitoes.
31	2	5"	
10/19/73	1	5 " `	Damp. Cooler (below 60 F)
**	2	5"	· .
**	_	5 "	
10/25/73	1	5 "	Temp. too low.
tl	2	5"	
11	*****	5#	
11/3/73	-	5"	Weather cool.
17	_	5"	
11/20/73	1	5"	Weather cool.
11		5"	

C. Box 75 Watt - white light

CATCH	TIME
3	2#
3	5"
4	15"
1	30 "
-	2"
4	5"
2	15"
3	30 [#]
2	5 "
1	5"
-	5"
	3 4 1 - 4 2 3

-page 5-

\mathbf{DATE}	CATCH	TIME	REMARKS
10/19/73	2	5"	Temperature probably too low for
††		5"	this intensity.
ŧŧ	****	5"	
10/25/73	_	5 "	Weather too cool.
11	***	5"	
11/3/73	-	5 "	Cold
11	_	5 " .	
11/20/73	· -	5"	Cold - These runs act as controls.
11	-	5"	

D. Box 15 Watt - white light

エノ	wauu	_	MITT 06	TTEIL
DAT	Έ		CATCH	TIME
10/9/	′7 3		2	2"
11			-	5"
11			****	15 "
11,				30"
10/1]	./73		_	2"
11			3	5"
71	•			15"
#1				30 "
10/15	3 /7 3			5"
Ħ			***	5"
10/19	73		l	5"
51			_	5"
10/25	5/73		-	5"
11				5"

No other responses were observed for this intensity of white light.

* All experiments made during the Fall were done behind LakeSide towards the ravine.

E. Funnel 15 Watt - ultraviolet

CATCH	TIME	REMARKS
. 2	2"	Weather cool. Windy.
. 2	5 "	•
l	15"	
-	30 "	
4	2"	Temp. in 70's F.
3	5 "	Overcast
5	15"	
2	30 "	
. 4	15".	Weather clear and warm.
2	15"	•
. 1	15#	
1 .	15"	•
l	15"	
	15"	
3	15"	
	15"	
2	15"	Partly cloudy. Warm -
· 1	15"	
3	15"	
-	15"	
· •••	15"	
1	15"	
_	15"	Cloudy and warm.
-	15"	
2	15 "	
2	15"	
+ 27 55	15!	
	15"	
-	15"	•
, 2	15 "	
•••	15"	
) ===	15"	
1	15"	
	2 2 1 - 4 3 5 2 4 2 1 1 1 - 3 - 2 1 3 1 - 2	2 2" 5 " 15" 30" 4 5" 5 " 5 " 5 " 15" 2 15" 1 15

DATE	CATCH	TIME	REMARKS
4/26/74	1	15"	with insecticide(commercial)
11 -	1	15"	
Ħ	-	15"	without insecticide
Ħ	_	15"	Weather damp. Partly cloudy,
11	-	15"	
4/28/74	1	15"	Used Arsenic compound as insect-
ŧŧ	***	15"	icide.
11		15"	Weather clear.
19	2	15"	
††	-	15"	

F. Funnel

15 Watt - white light (control)

DATE	CATCH	TIME	
3/27/74	1	2"	
11	3	5"	
***	3	15"	
†1	2	30 "	
4/5/74	3	2"	
11	2	5 "	
. 11	2	15"	
**	5	30 "	
4/6/74	3	155	
· £ ##	l	15"	
4/12/74	3	15"	
**	l	15"	
4/13/74	***	15"	
11	3	15"	
4/20/74	3	15"	
11	l	15"	
4/26/74	5	15"	With insecticide
11	2	15"	Without insecticide
4/28/74	1	15"	Used Arsenic compound
11	3	15"	

^{*} All experiments, using the funnel as the collector, were done behind Daniel-South (facing the Ouachita River).

V. ANALYSIS OF RESULTS

Although theoretically insects react more strongly to ultraviolet light than to ordinary white light, such was not the case in this experiment. There are several reasons for this deviation from the expected. One is the fact that there were other lights (within the range of the insects' vision) of greater intensity than those lights used in the experiment. This would cause the insects to react towards the more intense light source instead of the collector light. Although this reinforces the previously stated theory of "the greater the intensity, the greater the response", it voids the results concerning the insects reaction to ultraviolet light.

Ultraviolet light will draw more response than a white light of corresponding intensity, but there is a point at which the intensity will over-rule the response to the ultraviolet light.

Another factor that may have led to incorrect results is that of the intensity of the moon-light. On cloudy nights the reponse was greater than the response illicited on clear nights.

The temperature also had a very definite effect on the results. The lower the temperature the less the response. This relationship between temperature and response is an indirect one rather than a direct one. It is most likely that the temperature either killed or slowed down the insects, and did not actually effect the electro-chemical response.

In order to obtain valid results one would need to maintain strict controls on the temperature, wind currents, surroundings and humidity. The method of using the numbers of insects collected as the basis for experimental conclusions about the response of insects to variations of intensity and wave-length is not actually a valid method. A more precise method would be to record (if possible) the electrical responses of a single visual cell of the compound eye.

The results of this experiment reinforced the theory of 'the greater the intensity, the greater the response'. The results concerning the response to the ultraviolet light were voided due to the fact that surroundings were not controlled strictly enough.